

Table 5.14d
Unadjusted Patient Survival, Living Donor Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	11,177	99.5%	0.1%	11,177	98.5%	0.1%	34,288	91.0%	0.2%	56,493	77.1%	0.3%
Age at Tx												
<1 Year	6	83.3%	15.2%	6	83.3%	15.2%	15	86.7%	8.8%	35	94.3%	3.9%
1-5 Years	157	100.0%	0.0%	157	99.4%	0.6%	544	96.0%	1.0%	960	95.0%	0.8%
6-11 Years	127	100.0%	0.0%	127	100.0%	0.0%	490	98.0%	0.7%	957	94.8%	1.1%
12-17 Years	279	100.0%	0.0%	279	99.6%	0.4%	1,084	97.2%	0.6%	2,001	91.1%	1.0%
18-34 Years	2,119	99.8%	0.1%	2,119	99.2%	0.2%	6,844	95.8%	0.3%	12,416	88.6%	0.5%
35-49 Years	3,325	99.7%	0.1%	3,325	99.1%	0.2%	10,312	93.5%	0.3%	17,601	80.8%	0.5%
50-64 Years	3,948	99.3%	0.1%	3,948	98.1%	0.2%	11,621	88.1%	0.4%	17,890	65.3%	0.7%
65+ Years	1,216	99.1%	0.3%	1,216	96.6%	0.5%	3,378	77.6%	1.0%	4,633	42.1%	1.9%
Recipient Ethnicity/Race												
White	7,457	99.4%	0.1%	7,457	98.3%	0.1%	22,812	90.4%	0.3%	37,954	76.4%	0.4%
African American	1,632	99.7%	0.1%	1,632	98.7%	0.3%	5,238	90.0%	0.6%	8,567	73.3%	0.9%
Hispanic/Latino	1,495	99.6%	0.2%	1,495	99.1%	0.2%	4,473	94.2%	0.5%	7,152	83.8%	0.8%
Asian	460	99.8%	0.2%	460	99.3%	0.4%	1,376	95.0%	0.8%	2,239	85.0%	1.5%
Other/Multi-race	133	100.0%	0.0%	133	99.2%	0.7%	386	86.1%	2.7%	578	75.2%	3.3%
Unknown	0	-	-	0	-	-	3	66.7%	27.2%	3	+	+
Recipient Gender												
Female	4,268	99.6%	0.1%	4,268	98.8%	0.2%	13,632	91.4%	0.3%	22,868	77.7%	0.5%
Male	6,909	99.5%	0.1%	6,909	98.4%	0.2%	20,656	90.7%	0.3%	33,625	76.7%	0.4%
Recipient Blood Type												
O	5,023	99.5%	0.1%	5,023	98.6%	0.2%	15,572	91.5%	0.3%	25,635	78.0%	0.5%
A	4,277	99.6%	0.1%	4,277	98.4%	0.2%	12,964	90.3%	0.4%	21,405	75.8%	0.5%
B	1,440	99.5%	0.2%	1,440	98.5%	0.3%	4,466	90.9%	0.6%	7,337	78.0%	0.9%
AB	437	99.5%	0.3%	437	98.9%	0.5%	1,286	91.1%	1.1%	2,110	77.5%	1.7%
Unknown	0	-	-	0	-	-	0	-	-	6	100.0%	0.0%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14d (Continued)
Unadjusted Patient Survival, Living Donor Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	11,177	99.5%	0.1%	11,177	98.5%	0.1%	34,288	91.0%	0.2%	56,493	77.1%	0.3%
Recipient Residency												
U.S. Resident	11,089	99.5%	0.1%	11,089	98.5%	0.1%	33,982	90.9%	0.2%	55,898	77.0%	0.3%
Nonresident Alien	85	100.0%	0.0%	85	98.8%	1.2%	303	96.1%	1.2%	554	93.9%	1.2%
Unknown	3	100.0%	0.0%	3	100.0%	0.0%	3	+	+	41	75.5%	7.1%
Hospitalized at Tx												
Not Hospitalized	10,991	99.5%	0.1%	10,991	98.5%	0.1%	33,643	90.9%	0.2%	55,209	77.2%	0.3%
Hospitalized	158	98.7%	0.9%	158	96.8%	1.4%	557	91.9%	1.4%	1,114	75.8%	1.9%
In Intensive Care	22	95.5%	4.4%	22	95.5%	4.4%	82	88.1%	3.8%	151	69.1%	6.2%
Unknown	6	100.0%	0.0%	6	100.0%	0.0%	6	+	+	19	86.8%	8.9%
Primary Diagnosis												
Glomerular Diseases	3,151	99.7%	0.1%	3,151	99.1%	0.2%	9,886	95.1%	0.3%	16,936	86.5%	0.4%
Diabetes	2,436	99.4%	0.2%	2,436	97.9%	0.3%	7,498	83.5%	0.6%	12,196	58.4%	0.9%
Hypertensive												
Nephrosclerosis	1,918	99.3%	0.2%	1,918	98.0%	0.3%	5,141	89.0%	0.6%	7,691	72.2%	1.0%
Polycystic Kidneys	1,253	99.6%	0.2%	1,253	99.3%	0.2%	3,715	95.6%	0.5%	5,649	85.3%	0.9%
Tubular and Interstitial Diseases	613	99.3%	0.3%	613	98.4%	0.5%	2,126	89.8%	0.9%	3,814	79.5%	1.0%
Renovascular & Other Vascular Diseases	235	99.1%	0.6%	235	97.9%	0.9%	1,072	90.0%	1.1%	1,989	70.3%	1.9%
Congenital, Rare Familial, & Metabolic Disorders	377	99.7%	0.3%	377	98.7%	0.6%	1,329	96.5%	0.6%	2,258	90.0%	1.1%
Neoplasms	43	100.0%	0.0%	43	100.0%	0.0%	144	85.3%	3.9%	240	70.6%	6.0%
Other	1,132	99.6%	0.2%	1,132	98.3%	0.4%	3,324	92.0%	0.6%	5,646	81.4%	0.9%
Unknown	19	100.0%	0.0%	19	94.7%	5.1%	53	79.6%	8.1%	74	70.5%	8.9%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14d (Continued)
Unadjusted Patient Survival, Living Donor Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	11,177	99.5%	0.1%	11,177	98.5%	0.1%	34,288	91.0%	0.2%	56,493	77.1%	0.3%
PRA at Transplant												
0-9%	6,698	99.6%	0.1%	6,698	98.7%	0.1%	18,634	91.1%	0.3%	27,859	76.2%	0.5%
10-79%	845	99.3%	0.3%	845	98.8%	0.4%	1,975	89.3%	1.1%	2,643	71.1%	2.1%
80%+	115	99.1%	0.9%	115	96.5%	1.7%	285	87.5%	3.3%	358	77.7%	4.5%
Unknown	3,519	99.5%	0.1%	3,519	98.2%	0.2%	13,394	91.0%	0.3%	25,633	78.2%	0.4%
Yearly Center Volume												
0-12	179	99.4%	0.6%	179	99.4%	0.6%	556	94.2%	1.3%	1,429	84.0%	1.5%
13-28	507	99.0%	0.4%	507	99.0%	0.4%	1,808	91.5%	0.8%	3,786	78.4%	1.2%
29-47	955	99.1%	0.3%	955	98.1%	0.4%	3,117	89.5%	0.7%	5,859	75.5%	1.0%
48-81	1,827	99.5%	0.2%	1,827	98.8%	0.3%	6,657	90.8%	0.5%	12,135	76.2%	0.7%
82+	7,709	99.6%	0.1%	7,709	98.4%	0.1%	22,150	91.1%	0.3%	33,284	77.3%	0.4%
Dialysis Needed Within First Week After Tx												
No	10,744	99.6%	0.1%	10,744	98.6%	0.1%	32,815	91.5%	0.2%	53,830	77.9%	0.3%
Yes	426	97.7%	0.7%	426	95.8%	1.0%	1,466	79.1%	1.3%	2,562	62.4%	1.6%
Unknown	7	100.0%	0.0%	7	100.0%	0.0%	7	+	+	101	77.6%	4.3%
Donor Age												
6-11 Years	0	-	-	0	-	-	0	-	-	1	100.0%	0.0%
12-17 Years	3	100.0%	0.0%	3	100.0%	0.0%	4	100.0%	0.0%	11	100.0%	0.0%
18-34 Years	3,457	99.7%	0.1%	3,457	99.0%	0.2%	10,927	92.3%	0.3%	18,526	79.6%	0.5%
35-49 Years	5,044	99.5%	0.1%	5,044	98.6%	0.2%	15,948	91.0%	0.3%	26,352	77.3%	0.5%
50-64 Years	2,534	99.4%	0.2%	2,534	97.9%	0.3%	7,065	89.3%	0.5%	11,027	73.9%	0.8%
65+ Years	138	98.6%	1.0%	138	95.7%	1.7%	343	76.9%	3.2%	567	46.1%	3.9%
Unknown	1	0.0%	0.0%	1	0.0%	0.0%	1	0.0%	0.0%	9	77.8%	13.9%
Relation of Donor to Recipient												
Parent	1,125	99.9%	0.1%	1,125	99.7%	0.2%	3,891	96.1%	0.4%	7,465	89.1%	0.6%
Offspring	2,076	99.3%	0.2%	2,076	98.2%	0.3%	6,454	85.3%	0.6%	10,349	60.1%	1.0%
Sibling	2,891	99.6%	0.1%	2,891	98.7%	0.2%	9,486	93.6%	0.3%	17,410	82.6%	0.5%
Other Relative	857	99.9%	0.1%	857	98.6%	0.4%	2,590	91.2%	0.7%	4,121	78.0%	1.2%
Spouse	1,402	99.4%	0.2%	1,402	98.0%	0.4%	4,082	89.2%	0.7%	6,697	71.4%	1.1%
Other Unrelated	2,593	99.3%	0.2%	2,593	98.2%	0.3%	7,278	90.4%	0.5%	9,757	74.3%	1.2%
Unknown	233	99.6%	0.4%	233	99.1%	0.6%	507	88.1%	2.1%	694	77.8%	2.8%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14d (Continued)
Unadjusted Patient Survival, Living Donor Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	11,177	99.5%	0.1%	11,177	98.5%	0.1%	34,288	91.0%	0.2%	56,493	77.1%	0.3%
A Locus Mismatch												
0	2,473	99.5%	0.1%	2,473	98.5%	0.2%	8,063	92.2%	0.4%	14,263	79.9%	0.6%
1	5,981	99.5%	0.1%	5,981	98.6%	0.2%	18,521	90.4%	0.3%	30,451	76.4%	0.4%
2	2,610	99.5%	0.1%	2,610	98.3%	0.3%	7,423	91.0%	0.5%	11,018	75.1%	0.9%
Unknown	113	98.2%	1.2%	113	98.2%	1.2%	281	88.8%	2.5%	761	75.8%	2.1%
B Locus Mismatch												
0	1,687	99.6%	0.2%	1,687	98.8%	0.3%	5,579	93.0%	0.5%	10,164	81.5%	0.6%
1	5,764	99.5%	0.1%	5,764	98.6%	0.2%	17,962	90.6%	0.3%	29,966	76.6%	0.4%
2	3,613	99.5%	0.1%	3,613	98.3%	0.2%	10,466	90.4%	0.4%	15,603	74.7%	0.7%
Unknown	113	98.2%	1.2%	113	98.2%	1.2%	281	88.8%	2.5%	760	75.8%	2.1%
DR Locus Mismatch												
0	2,359	99.8%	0.1%	2,359	98.9%	0.2%	7,547	92.3%	0.4%	13,352	80.5%	0.6%
1	5,937	99.4%	0.1%	5,937	98.5%	0.2%	18,489	90.9%	0.3%	30,604	76.3%	0.4%
2	2,769	99.5%	0.1%	2,769	98.3%	0.2%	7,966	89.8%	0.5%	11,730	74.8%	0.9%
Unknown	112	98.2%	1.3%	112	98.2%	1.3%	286	89.2%	2.4%	807	76.4%	2.0%
Total Mismatch												
0	911	99.7%	0.2%	911	98.8%	0.4%	2,980	94.4%	0.6%	5,587	84.3%	0.8%
1	597	99.7%	0.2%	597	98.2%	0.6%	2,030	90.9%	0.8%	3,596	78.9%	1.2%
2	1,910	99.6%	0.1%	1,910	99.0%	0.2%	6,202	91.1%	0.5%	10,765	76.4%	0.7%
3	3,209	99.5%	0.1%	3,209	98.6%	0.2%	10,042	90.4%	0.4%	16,690	76.4%	0.6%
4	1,586	99.2%	0.2%	1,586	98.0%	0.3%	4,522	91.0%	0.6%	6,938	76.9%	1.0%
5	1,827	99.7%	0.1%	1,827	98.4%	0.3%	5,326	90.1%	0.6%	7,953	74.3%	1.0%
6	1,023	99.4%	0.2%	1,023	98.2%	0.4%	2,897	90.3%	0.8%	4,151	74.0%	1.5%
Unknown	114	98.2%	1.2%	114	98.2%	1.2%	289	89.3%	2.4%	813	76.1%	2.0%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14d (Continued)
Unadjusted Patient Survival, Living Donor Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	11,177	99.5%	0.1%	11,177	98.5%	0.1%	34,288	91.0%	0.2%	56,493	77.1%	0.3%
Transplant Center State												
Alabama	280	100.0%	0.0%	280	100.0%	0.0%	896	90.1%	1.4%	1,446	75.0%	2.0%
Arkansas	48	100.0%	0.0%	48	97.9%	2.1%	192	90.1%	2.7%	389	67.3%	4.0%
Arizona	304	99.7%	0.3%	304	99.0%	0.6%	811	90.2%	1.4%	1,198	76.8%	2.4%
California	1,202	99.8%	0.1%	1,202	99.0%	0.3%	3,724	93.0%	0.6%	6,256	80.9%	0.9%
Colorado	198	99.5%	0.5%	198	98.5%	0.9%	510	91.9%	1.6%	841	76.4%	2.8%
Connecticut	107	100.0%	0.0%	107	99.1%	0.9%	297	92.7%	2.0%	623	78.6%	2.6%
District of Columbia	157	100.0%	0.0%	157	100.0%	0.0%	518	93.1%	1.4%	954	74.1%	2.5%
Delaware	9	100.0%	0.0%	9	100.0%	0.0%	18	94.4%	5.4%	24	95.8%	4.1%
Florida	381	99.2%	0.5%	381	98.2%	0.7%	1,119	88.7%	1.4%	1,815	75.3%	1.8%
Georgia	216	100.0%	0.0%	216	99.1%	0.7%	661	92.2%	1.4%	1,026	79.8%	2.4%
Hawaii	25	96.0%	3.9%	25	96.0%	3.9%	105	90.5%	3.6%	150	83.3%	4.4%
Iowa	65	100.0%	0.0%	65	98.5%	1.5%	257	89.6%	2.5%	472	75.9%	3.2%
Illinois	605	99.5%	0.3%	605	98.5%	0.5%	1,790	90.8%	0.9%	2,737	78.8%	1.5%
Indiana	176	100.0%	0.0%	176	98.9%	0.8%	433	96.5%	1.2%	687	85.3%	2.3%
Kansas	37	100.0%	0.0%	37	100.0%	0.0%	124	94.0%	2.8%	221	81.0%	4.2%
Kentucky	64	98.4%	1.6%	64	98.4%	1.6%	203	91.2%	2.8%	360	86.8%	2.5%
Louisiana	135	99.3%	0.7%	135	97.8%	1.3%	434	90.9%	1.8%	701	76.9%	2.7%
Massachusetts	367	99.5%	0.4%	367	99.2%	0.5%	1,067	92.1%	1.2%	1,814	80.2%	1.6%
Maryland	299	99.3%	0.5%	299	97.0%	1.0%	922	91.2%	1.2%	1,820	73.6%	1.6%
Maine	47	100.0%	0.0%	47	97.9%	2.1%	139	91.7%	2.8%	235	76.8%	5.2%
Michigan	425	99.3%	0.4%	425	97.6%	0.7%	1,411	88.4%	1.1%	2,407	73.9%	1.5%
Minnesota	579	99.1%	0.4%	579	97.9%	0.6%	1,843	89.6%	0.9%	3,003	75.1%	1.4%
Missouri	239	99.6%	0.4%	239	98.7%	0.7%	666	93.7%	1.3%	1,198	77.2%	2.2%
Mississippi	3	100.0%	0.0%	3	100.0%	0.0%	8	100.0%	0.0%	24	79.6%	9.2%
North Carolina	215	99.1%	0.7%	215	97.2%	1.1%	813	88.6%	1.4%	1,418	78.2%	1.8%
North Dakota	41	100.0%	0.0%	41	100.0%	0.0%	146	91.1%	3.2%	230	71.8%	5.8%
Nebraska	112	99.1%	0.9%	112	99.1%	0.9%	281	94.9%	1.7%	410	82.3%	4.2%
New Hampshire	37	100.0%	0.0%	37	100.0%	0.0%	96	95.5%	3.1%	143	83.4%	5.4%
New Jersey	329	99.4%	0.4%	329	98.5%	0.7%	1,009	92.0%	1.2%	1,537	77.6%	2.1%
New Mexico	35	100.0%	0.0%	35	100.0%	0.0%	121	88.4%	4.2%	216	82.4%	4.0%
Nevada	22	100.0%	0.0%	22	100.0%	0.0%	83	89.1%	4.3%	164	69.7%	6.8%
New York	938	99.5%	0.2%	938	98.5%	0.4%	2,658	92.3%	0.7%	4,250	78.1%	1.2%
Ohio	644	99.7%	0.2%	644	97.8%	0.6%	1,954	88.6%	1.0%	2,948	70.7%	1.8%
Oklahoma	39	97.4%	2.5%	39	94.9%	3.5%	167	88.1%	3.0%	289	76.3%	4.1%
Oregon	150	99.3%	0.7%	150	98.7%	0.9%	536	92.1%	1.6%	856	79.8%	2.6%

(Continued)

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.

Table 5.14d (Continued)
Unadjusted Patient Survival, Living Donor Kidney Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months			1 Year			5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Transplant Center State												
Pennsylvania	565	99.3%	0.4%	565	98.6%	0.5%	1,779	91.3%	0.9%	2,745	78.2%	1.4%
Puerto Rico	21	100.0%	0.0%	21	100.0%	0.0%	114	91.2%	3.1%	253	72.0%	4.5%
Rhode Island	48	100.0%	0.0%	48	100.0%	0.0%	167	91.9%	2.3%	290	69.4%	5.3%
South Carolina	63	98.4%	1.6%	63	98.4%	1.6%	157	91.3%	2.8%	329	73.4%	3.6%
South Dakota	29	100.0%	0.0%	29	93.1%	4.7%	100	83.0%	5.2%	123	79.1%	5.6%
Tennessee	245	99.6%	0.4%	245	97.1%	1.1%	802	88.3%	1.5%	1,309	75.0%	2.0%
Texas	649	99.2%	0.3%	649	98.5%	0.5%	1,931	90.4%	0.9%	3,154	77.8%	1.3%
Utah	171	100.0%	0.0%	171	99.4%	0.6%	457	94.1%	1.5%	771	82.4%	2.2%
Virginia	327	99.4%	0.4%	327	98.8%	0.6%	1,032	88.0%	1.4%	1,632	75.7%	1.9%
Vermont	27	96.3%	3.6%	27	92.6%	5.0%	59	94.9%	2.9%	93	86.6%	6.6%
Washington	201	100.0%	0.0%	201	100.0%	0.0%	696	89.7%	1.5%	1,269	78.6%	1.7%
Wisconsin	279	99.6%	0.4%	279	98.9%	0.6%	891	92.2%	1.2%	1,488	78.1%	1.8%
West Virginia	22	100.0%	0.0%	22	100.0%	0.0%	91	88.1%	5.1%	175	65.5%	6.6%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Dialysis needed within first week excludes patients whose graft failed before the end of the week.